

## Patient Information

# FLOXIN® (FLOX-IN) Otic SINGLES®

(ofloxacin otic) solution 0.3%

### IMPORTANT PATIENT INFORMATION AND INSTRUCTIONS. READ BEFORE USE.

**THROW PLASTIC DISPENSING CONTAINER AWAY IMMEDIATELY AFTER USE. THE CONTAINER AND TOP, DUE TO THEIR SMALL SIZE, POSE A DANGER OF CHOKING TO YOUNG CHILDREN.**

#### What is FLOXIN Otic SINGLES®?

FLOXIN Otic SINGLES® is an antibiotic in a sterile solution used to treat ear infections caused by certain bacteria found in:

- patients (12 years and older) who have a middle ear infection and have a hole in the eardrum
- pediatric patients (between 1 and 12 years of age) who have a middle ear infection and have a tube in the eardrum
- patients (6 months and older) who have an infection in the ear canal

**Middle Ear Infection:** A middle ear infection is a bacterial infection behind the eardrum. People with a hole or a tube in the eardrum may notice a discharge (fluid draining) from the ear canal.

**Ear Canal Infection:** An ear canal infection (also known as "Swimmer's Ear") is a bacterial infection of the ear canal. The ear canal and the outer part of the ear may swell, turn red, and be painful. Also, a fluid discharge may appear in the ear canal.


#### Who should NOT use FLOXIN Otic SINGLES®?

- Do not use this product if you are allergic to ofloxacin or to other quinolone antibiotics
- Do not give this product to pediatric patients who:
  - have an ear canal infection and are less than 6 months of age because no data were collected from this population
  - have a middle ear infection and have a tube in the eardrum and are less than one year of age because no data were collected from this population
  - have a middle ear infection and have a hole in the eardrum and are less than 12 years of age because no data were collected from this population

#### How should FLOXIN Otic SINGLES® be given?

##### 1. Wash hands

The person giving FLOXIN Otic SINGLES® should wash his/her hands with soap and water.



##### 2. Clean ear & warm single-dispensing container

Gently clean any discharge that can be removed easily from the outer ear. DO NOT INSERT ANY OBJECT OR SWAB INTO THE EAR CANAL.

Hold the container of FLOXIN Otic SINGLES® in the hand for one or two minutes to warm the solution.



##### 3. Apply contents of container(s)

Tap container to make the solution settle at the bottom of the container. Twist the container top to open and discard top immediately.

###### For a Middle Ear Infection:

The person receiving FLOXIN Otic SINGLES® should lie on his/her side with the infected ear up. Patients (12 and older) apply the contents of **2 single-dispensing containers** of FLOXIN Otic SINGLES® into the infected ear.

For pediatric patients under 12 years of age, apply the contents of

**1 single-dispensing container** into the infected ear. The tip of the container should not touch the fingers or the ear or any other surfaces.

###### For an Ear Canal Infection ("Swimmer's Ear"):

The person receiving FLOXIN Otic SINGLES® should lie on his/her side with the infected ear up. Patients (13 and older) apply the contents of **2 single-dispensing containers** of FLOXIN Otic SINGLES® into the infected ear. For pediatric patients under 13 years of age, apply the contents of **1 single-dispensing container** into the infected ear. The tip of the container should not touch the fingers or the ear or any other surfaces.

**BE SURE TO FOLLOW THE INSTRUCTIONS BELOW FOR THE PATIENT'S SPECIFIC EAR INFECTION.**

##### 4. Press ear or pull ear

###### For a Middle Ear Infection:

While the person receiving FLOXIN Otic SINGLES® lies on his/her side, the person giving the medication should gently press the **TRAGUS** (see diagram) 4 times in a pumping motion. This will allow the drops to pass through the hole or tube in the eardrum and into the middle ear.



###### For an Ear Canal Infection ("Swimmer's Ear"):

While the person receiving the medication lies on his/her side, the person giving the drops should gently pull the outer ear upward and backward. This will allow the medication to flow down into the ear canal.

TRAGUS



##### 5. Stay on side

The person who received the medication should remain on his/her side for at least 5 minutes.



**Repeat steps 2-5 for the other ear if both ears are infected.**

#### How often should FLOXIN Otic SINGLES® be given?

In patients with an Ear Canal Infection ("Swimmer's Ear"), FLOXIN Otic SINGLES® should be given once daily at about the same time each day (for example, 8 AM or 8 PM) in each infected ear unless the doctor has instructed otherwise.

In patients with a Middle Ear Infection, FLOXIN Otic SINGLES® should be given 2 times each day (about 12 hours apart, for example 8 AM and 8 PM) in each infected ear unless the doctor has instructed otherwise. The best times to use the medication are in the morning and at night.

It is very important to use the medication for as long as the doctor has instructed, **even if the symptoms improve**. If FLOXIN Otic SINGLES® is not used for as long as the doctor has instructed, the infection may be more likely to return.

#### What if a dose is missed?

In patients with an Ear Canal Infection ("Swimmer's Ear"), it is important that you take the medication every day. If you miss a dose which may have been scheduled for earlier in the day (for example, 8 AM), you should take that day's dose as soon as possible and then go back to your regular daily dosing schedule.

In patients with a Middle Ear Infection, if a dose of FLOXIN Otic SINGLES® is missed, it should be given as soon as possible. However, if it is almost time for the next dose, skip the missed dose and go back to the regular dosing schedule.

Do not use a double dose unless the doctor has instructed you to do so. If the infection is not improved after one week, you should consult your doctor. If you have two or more episodes of drainage within six months, it is recommended that you see your doctor for further evaluation.

#### What activities should be avoided while using FLOXIN Otic SINGLES®?

It is important that the infected ear(s) remain clean and dry. When bathing, avoid getting the infected ear(s) wet. Avoid swimming unless the doctor has instructed otherwise.

#### What are some of the possible side effects of FLOXIN Otic SINGLES®?

During the testing of FLOXIN Otic SINGLES® in external ear infections, the most common side effect was discomfort upon application which happened in 7% of patients. If the pain is severe, the medication should be stopped and you should contact your doctor. Other side effects were: itching (1%), earache (0.8%), and dizziness (0.4%).

During testing of FLOXIN Otic SINGLES® in middle ear infections, the most common side effect was a bitter taste which happened in 7% of patients with a middle ear infection. This may occur when some of the medication passes from the middle ear to the back of the mouth. This side effect is not serious and there is no need to stop the medicine if this should happen. Other side effects which were found in 1% of the patients were: earache, itching, abnormal sensation, rash and dizziness.

Call your doctor about these or other side effects if they occur.

If a rash or an allergic reaction to FLOXIN Otic SINGLES® occurs, stop using the product and contact your doctor.

#### DO NOT TAKE FLOXIN Otic SINGLES® BY MOUTH.

If FLOXIN Otic SINGLES® is accidentally swallowed or an overdose occurs, call the doctor immediately. This medicine is available only with a doctor's prescription. Use only as directed. Do not use this medicine if outdated.

If you wish to learn more about FLOXIN Otic SINGLES®, ask the doctor or pharmacist. Complete prescribing information is printed on the reverse side.

#### HOW SUPPLIED

Single-dispensing containers, 0.25 mL each, 2 per foil pouch.

**Storage Conditions:** Store at 25°C (77°F), excursions permitted to 15–30°C (59–86°F).

**Store containers in pouch to protect from light.**

#### FLOXIN Otic SINGLES® is manufactured for:

Daiichi Pharmaceutical Corporation  
Montvale, NJ 07645 USA

This Patient Information has been approved by the U.S. Food and Drug Administration.

Revised: April 2005

Copyright© 2004 Daiichi Pharmaceutical Corporation. All rights reserved.

Covered by U.S. Patent No. 5,401,741